


	L #	Hits	Search Text	DBs	Time Stamp
1	L1	8	inositol adj phosphoglycan	USPA T	2003/12/1 2 16:14
2	L2	13	(2f7 or 2d1 or 5h6) and 530/387.1,388.1.ccls.	USPA T	2003/12/1 2 16:15
3	L3	70092	cancer or tumor	USPA T	2003/12/1 2 16:15
4	L4	22	gpi-anchor\$3 and ((cancer or tumor or proliferat\$3 or growth).clm.)	USPA T	2003/12/1 2 16:15
5	L5	6	inositolphosphoglycan	USPA T	2003/12/1 2 16:15
6	L6	662	ipg	USPA T	2003/12/1 2 16:15
7	L7	32	3 and 6	USPA T	2003/12/1 2 16:15



Entrez PubMed

Nucleotide

Protein

Genome

Structure

PMC

Journals

Bo

Search PubMed

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show:

20

Sort

Send to Text

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Cell Signal. 1995 May;7(4):411-21.

Related Articles, L

ELSEVIER  
FULL-TEXT ARTICLE**Role of glycosyl-phosphatidylinositol hydrolysis as a mitogenic signal for epidermal growth factor.****Clemente R, Jones DR, Ochoa P, Romero G, Mato JM, Varela-Nieto I.**

Instituto de Investigaciones Biomedicas, Consejo Superior de Investigaciones Cientificas (CSIC), Madrid, Spain.

We have investigated the role of the hydrolysis of glycosyl-phosphatidylinositol (GPI) as one of the signalling pathways elicited after interaction of epidermal growth factor (EGF) with its specific plasma membrane receptor (EGFR). Endogenous GPI was characterized in both NIH 3T3 mouse fibroblast cells and in EGFR-transfected NIH 3T3 cells (designated EGFR T17). GPI molecules isolated from both cell lines were identical and they incorporated radioactivity from both sugar and fatty acid substrates. Incubation of EGFR T17 cells with EGF, produced a rapid and transient hydrolysis of GPI. Maximum hydrolysis occurred after a 1-min incubation with 50 nM EGF. No such effects of EGF were observed in the parental cell line. Both inositol phosphoglycan (IPG)- and EGF-induced cell proliferation was inhibited in the presence of an IPG-antibody to different extents. The relationship between GPI hydrolysis and the activity of the EGFR was studied using the tyrosine kinase inhibitors tyrphostin (RG50864) and genistein. These agents were able to significantly inhibit EGF-mediated cell proliferation, EGF-dependent hydrolysis of GPI and EGF-regulated autophosphorylation of the EGFR. It is concluded that GPI hydrolysis is one of the earliest intracellular events generated in response to EGF.

PMID: 8527310 [PubMed - indexed for MEDLINE]

Display

Abstract

Show:

20

Sort

Send to Text

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act](#) | [Disclaimer](#)

Dec 1 2003 07: